

Peace Harbor Medical Center Outpatient Infusion Service Request

Phone: (541) 902-6019 Fax: (541) 902-1649

Thank you for selecting our infusion services team to care for your patient. If you are not a PeaceHealth provider, all information listed below is required before we can process orders and schedule your patient for treatment. Part A- Patient scheduling and contact information: Patient Name (Last, First): _______Date of Birth: _____ Patient Contact Information and Phone Number (s): Ordering Provider Name (Print): Provider Clinic or Service Address: ______ Clinic or Service Phone Number: _____ Clinic or Service Fax Number: _____ Diagnosis (include ICD 10 codes): Medication and Service Requested- list J-Code/ CPT code if known: ______ Date Service is Requested to Begin: ______ Date Service is Expected to End: ___ Order will expire 1 year from date of provider signature unless "date service is expected to end" is earlier. <u>Part B- Insurance and Prior Authorization.</u> Any non-PeaceHealth provider must obtain prior authorization prior to service. Attach a copy of authorization documentation received from insurance payer when submitting orders. Insurance (Payer) Company: Prior Authorization Number and Conditions: Prior Authorization Expiration Date: _____ Insurance (Payer) Contact Phone Number: <u>Part C-</u> Elements needed to guide medication therapy are included with request for service: All orders and instruction (please use the PeaceHealth approved ordering form) are complete and include provider signature AND printed name at the bottom of each order page. Check the boxes of ALL orders you would like to activate. For blood products, PeaceHealth Blood and Transfusion Consent form is signed and dated by the provider and the patient. If information is located outside of PeaceHealth's electronic medical record system attach the following: A list of current medications reconciled by patient provider is available and includes a list of known allergies. Recent progress notes from ordering provider. A copy of relevant laboratory results and other appropriate supporting documentation. **IMPORTANT MESSAGE TO PROVIDERS:** To reduce delays in treatment and phone calls to your office you may participate in the PHMC formulary process by signing this document. A clinical pharmacist will adjust orders according to PHMC approved policies and procedures.

I agree to utilize PHMC policies & procedures that have been reviewed by the Pharmacy & Therapeutics Committee and authorized by the Medical Executive Committee of PHMC. This agreement will be issued for the duration of active orders contained within this treatment plan.

PROVIDER SIGNATURE: ______ DATE: _____ TIME:_____

FAX completed service request and completed orders to: PHMC OP Infusion and Nursing Services 541-902-1649



Progress & Orders



Magnesium **Outpatient Infusion Therapy Plan**

Heading	ders Are Initiated by Default Unless Crossed Out by Practitioner. All <u>Boxed Orders</u> Require Practitioner Check to be Initiated. Content		
Supportive Care	Choose one of the following:		
	Magnesium sulfate gram IV once		
	☐ Magnesium sulfate IV weekly based on lab value. Check lab results prior to releasing this order:		
	For magnesium serum level 1.5-1.9 mg/dL, give 2 grams magnesium sulfate IV		
	 For magnesium serum level 0.9-1.4 mg/dL, give 4 grams magnesium sulfate IV 		
	For magnesium serum level less than 0.9 mg/dL, start 4 grams magnesium sulfate IV and call		
	provider for additional orders		
Labs	Serum magnesium level prior to each treatment		
	☐ Treatment lab instructions – Provider approves to release and draw labs 2 days pre and post this planned treatment date.		
	·		
Nursing Orders	□ Nursing communication – Check serum magnesium weekly and infuse as follows for lab value:		
	1.5-1.9 mg/dL: give 2 grams magnesium sulfate IV		
	0.9-1.4 mg/dL: give 4 grams magnesium sulfate IV		
	 Less than 0.9 mg/dL: start 4 grams magnesium sulfate IV and call provider for added orders. 		
Nursing IV Access	Select the most appropriate option below:		
and Maintenance	☐ Insert PERIPHERAL IV as needed and flush (unless provider selects option for a central line).		
	☑ Sodium chloride 0.9% (NS) flush 10 mL IV once as needed for line care.		
	Access and use NON DICC Control Line (CVAD		
	☐ Access and use NON-PICC Central Line/CVAD		
	☐ Initiate Central Line (non-PICC) maintenance protocol.		
	Sodium chloride 0.9% (NS) flush 10 mL IV as needed for line care, before and after medication		
	administration, at discharge, and at de-access (sterile NS for Port-a-Cath access)		
	☑ Sodium chloride 0.9% (NS) flush 20 mL IV as needed for line care post lab draw.		
	☐ Heparin, porcine (PF) 100 unit/mL flush 5 mL IV as needed for line care, for de-access.		
	☐ Alteplase (Cathflo) injection 2 mg intra-catheter as needed for occluded central line catheters-		
	For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand		
	undisturbed to allow large bubbles to dissipate. Mix by gently swirling until completely		
	dissolved (complete dissolution should occur within 3 minutes); do not shake. Final		
	concentration: 1mg/mL. Retain in catheter for 30 minutes to 2 hours; may instill a second dose		
	if occluded.		
	Access and use DICC Control Line (CVAD		
	☐ Access and use PICC Central Line/CVAD		
	☐ Initiate PICC maintenance protocol.		
	☐ Change PICC line dressing weekly and as needed.		
	☑ Sodium chloride 0.9% (NS) flush 10 mL IV as needed for line care, and before and after		
	medication administration.		
	Sodium chloride 0.9% (NS) flush 20 mL IV as needed for line care post lab draw		
	☐ Alteplase (Cathflo) injection 2 mg intra-catheter as needed for occluded central line catheters-		
	For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand		
	undisturbed to allow large bubbles to dissipate. Mix by gently swirling until completely		
	dissolved (complete dissolution should occur within 3 minutes); do not shake. Final		
	1		

Practitioner Signature:	Date of Order: _	Time:

Final page of orders must include signature of the ordering practitioner, date, and time.



Progress & Orders



Magnesium Outpatient Infusion Therapy Plan

All Pre-Selected Boxed Orders Are Initiated by Default Unless Crossed Out by Practitioner. All Boxed Orders Require Practitioner Check to be Initiated.

Heading	Content		
	concentration: 1mg/mL. Retain in catheter for 30 minutes to 2 hours; may instill a second dose		
	if occluded.		
As Needed	☑ DiphenhydrAMINE (Benadryl) injection 50 mg IV as needed for itching or hives. May give an		
Medications	additional 50 mg PO or IV 4 hours after first dose.		
	☐ DiphenhydrAMINE (Benadryl) 50 mg PO as needed for itching or hives. May give an additional 50		
	mg PO or IV 4 hours after first dose.		
	☐ Sodium chloride 0.9% (NS) flush 10 mL IV as needed for line care.		
	☑ Sodium chloride 0.9% 500 mL continuous infusion at 25 mL/hour IV as needed for therapy		
	administration (i.e., blood products, chemotherapy, potassium administration).		
Referral	☐ Ambulatory referral to OP Infusion Services		
PHMC Outpatient	PROVIDER – PLEASE SIGN, DATE AND TIME ORDERS AND RETURN TO:		
Infusion Contact	PeaceHealth Peace Harbor Medical Center Outpatient Infusion Services Department		
Information	400 Ninth Street, Florence, OR 97439		
	Contact Phone: 541-902-6019 and FAX 541-902-1649		
Authorization by	Person giving verbal or telephone order:		
Verbal or	Person receiving verbal or telephone order:		
Telephone Order	☐ Check to indicate verbal or telephone orders have been read back to confirm accuracy		

Practitioner Signature:	Date of Order:	Time: