

#### **SAVE A LIFE**

## STOP THE BLEED® Course American College of Surgeons

Copyright © 2019 American College of Surgeons

Version 2



BLEEDINGCONTROL.ORG
STOPTHEBLEED.ORG





#### Stop the Bleed® is a collaborative effort by:



The Hartford Consensus Group



The American
College of Surgeons
Committee on
Trauma



The Committee on Tactical Combat Casualty Care



The National
Association of
Emergency Medical
Technicians

WARNING! Some of the images shown during this presentation are graphic and may be disturbing to some people.

#### Why Do I Need This Training?

The #1 cause of preventable death after injury is bleeding.

#### Where Can I Use This Training?



#### Goals

#### 1. Identify

Recognize
Iife-threatening
bleeding

#### 2. Stop the Bleed

Take steps to STOP THE BLEEDING

- ✓ Pressure
- ✓ Packing
- ✓ Tourniquets

#### **Personal Safety**

#### YOUR safety is YOUR first priority

- If you are injured, you cannot help others
- Help others only when it's safe to do so
- If the situation changes or becomes unsafe:
  - √ Stop
  - ✓ Move to safety
  - ✓ If you can, take the victim with you

#### **Personal Safety**

#### YOUR safety is YOUR first priority


- Wear gloves if you can
- If you get blood on you, be sure to clean any part of your body that the blood has touched
- Tell a health care provider that you got blood on you, and follow his or her direction

A Alert 911

**B** Bleeding

C Compress

- A Alert 911
- Bleeding
- Compress



## A Alert 911

- Call 911
- Know your location
- Follow instructions provided by 911 operator

A Alert 911

**B** Bleeding

Compress



## **B** Bleeding


- Find source of bleeding
- Look for:
  - Continuous bleeding
  - Large-volume bleeding
  - Pooling of blood


## **B** Bleeding


- There may be multiple places the victim is bleeding
- Clothing may also hide life-threatening bleeding

## **B** Bleeding

- Arms and legs
- Neck, armpits, and groin
- Body







A Alert 911


Bleeding

C Compress - Pressure




## C Compress - Pressure


- Apply direct pressure to wound
- Focus on the location of the bleeding
- Use just enough gauze or cloth to cover injury
- If pressure stops the bleeding, keep pressure on wound until help arrives



# C Compress - Packing


- For large wounds, superficial pressure is not effective
- If bleeding is from a deep wound, pack gauze tightly into the wound until it stops the bleeding; hold pressure until help arrives







# C Compress - Packing

- Arms and legs
- Neck, armpits, and groin
- Body








# C Compress - Tourniquet

- Apply 2 to 3 inches above wound
- Do not place over the elbow or knee
- Tighten tourniquet until bleeding stops
- Do NOT remove the tourniquet

# C Compress - Tourniquet

- Can apply to others or on yourself
- Can be applied over clothes
- Tourniquets HURT
- A second tourniquet may be required to stop the bleeding





## Bleeding control in children

- In all but the extremely young child, the same tourniquet used for adults can be used in children.
- For the infant or very small child (tourniquet too big), direct pressure on the wound as described previously will work in virtually all cases.
- For large, deep wounds, wound packing can be performed in children just as in adults using the same technique as described previously.

#### **FAQs**

- Impaled objects?
- Improvised tourniquets?
- Loss of arm or leg?
- Pain?
- Other questions?

#### Summary

- ✓ Personal safety
- A Alert 911
- **B** Find bleeding
- C Compress with pressure and/or packing
- C Compress with a tourniquet
- ✓ Wait for help to arrive









#### For more information:

**BLEEDINGCONTROL.ORG** 

STOPTHEBLEED.ORG



The only thing more tragic than a death... is a death that could have been prevented.