

Height _____ Weight ____

Riverbend Medical Center Regional Infusion Center (RIC) 3377 Riverbend Drive Suite 502/510 Springfield, Oregon 97477 Phone 541-222-6280 Fax 541-349-8006

Denosumab (PROLIA) INJECTION (v. 05/16/2024)

Diagnos	sis/Indication (ICD-10):
	needs to have had satisfactory dental exam prior to start of denosumab (Prolia). Please send copy of most recent calcium h this order (must be within last 60 days).
Medication	as:
	Denosumab 60 mg subcutaneous injection every 6 months
Nursing co	mmunications:
•	Instruct patient to take calcium 1000 mg orally daily and at least 500 IU vitamin D daily Remind patient of good dental hygiene and to avoid dental procedures other than cleaning
Emergency	y Medications:
■ diapho	DiphenhydrAMINE (BENADRYL) 25 to 50 mg IV as needed for mild to moderate drug reactions (flushing, dizziness, headaches, presis, fever, palpitations, chest discomfort, blood pressure changes (>/= 20 points in SBP), nausea, urticaria, chills, pruritic).
mild to 20 poi	Administer 25 mg IV once, if reaction does not resolve in 3 minutes may repeat 25 mg IV dose for a total of 50 mg and contact provider. MethylPREDNISolone sodium succinate (Solu-MEDROL) 125 mg IV once as needed for shortness of breath, continued symptoms of a moderate drug reaction (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>/ onts in SBP), nausea, urticaria, chills, pruritic) that worsen or persist 5 minutes after administration of diphenhydramine (Benadryl). ct provider if given.
■ Avoid	Epinephrine 0.3 mg IM once for anaphylaxis. If reaction does not resolve in 3 minutes may repeat 0.3 mg IM dose for a total of 0.6 n use of hand, foot, leg veins in elderly patient and those with occlusive vascular disease. Contact provider if given.
	Famotidine (PEPCID) 20 mg IV once as needed for infusion/allergic reaction.
tient name: _	Provider printed name:
)B:	Provider signature:

Date: _____ Time: ____