



Riverbend Medical Center Regional Infusion Center (RIC)  
3377 Riverbend Drive Suite 502/510  
Springfield, Oregon 97477  
Phone 541-222-6280 Fax 541-349-8006

**PEGFILGRASTIM-cbqv (Udenyca) Injection (v. 01/11/2024)**

Diagnosis/Indication (ICD-10): \_\_\_\_\_

**Medication:**

■ Pegfilgrastim-cbqv 6 mg subcutaneous every \_\_\_\_\_ days for \_\_\_\_\_ doses

**Nursing communications:**

■ Vital signs: as needed

**Emergency Medications:**

- DiphenhydrAMINE (BENADRYL) 25 to 50 mg IV as needed for mild to moderate drug reactions (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes ( $\geq 20$  points in SBP), nausea, urticaria, chills, pruritic).
  - Administer 25 mg IV once, if reaction does not resolve in 3 minutes may repeat 25 mg IV dose for a total of 50 mg and contact provider.
- MethylPREDNISolone sodium succinate (Solu-MEDROL) 125 mg IV once as needed for shortness of breath, continued symptoms of mild to moderate drug reaction (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes ( $\geq 20$  points in SBP), nausea, urticaria, chills, pruritic) that worsen or persist 5 minutes after administration of diphenhydramine (Benadryl). Contact provider if given.
- Epinephrine 0.3 mg IM once for anaphylaxis. If reaction does not resolve in 3 minutes may repeat 0.3 mg IM dose for a total of 0.6 mg. Avoid use of hand, foot, leg veins in elderly patient and those with occlusive vascular disease. Contact provider if given.
- Famotidine (PEPCID) 20 mg IV once as needed for infusion/ allergic reaction.
- Emergency medications can be given IM if IV route unavailable

Patient name and DOB \_\_\_\_\_

Provider printed name: \_\_\_\_\_

Provider signature: \_\_\_\_\_

Height \_\_\_\_\_ Weight \_\_\_\_\_

Date: \_\_\_\_\_ Time: \_\_\_\_\_