

Riverbend Medical Center Regional Infusion Center (RIC) 3377 Riverbend Drive Suite 502/510 Springfield, Oregon 97477 Phone 541-222-6280 Fax 541-349-8006

Methylprednisolone (Solu-Medrol) Order Set (v. 05/21/2024)

Diagno	sis/Indication (ICD-10):		
Medicatio	ous:		
	Methylprednisolone 1 gram IVPB daily over 1 hour x days		
	Methylprednisolonemg IVPB over 1 hour daily x days		
Nursingco	ommunications:		
•	Vital signs: Initial and as needed during infusion		
Access:			
	Insert peripheral IV		
	 Every visit, remove after IV administration compl 	lete	
	Access & Use Central Line/ CVAD		
	 Initiate Central Line (Non-PICC) Maintenance Protocol Heparin, porcine (PF) 100 unit/mL flush 5 mL as needed for Port-a-Cath line care 		
	 Alteplase (Cathflo) 2 mg as needed for occluded catheter. For clearing central line catheter- retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded 		
	Access & Use PICC		
	- Initiate PICC Maintenance Protocol		
	- Normal saline flush 3 mL as needed for PICC/ Hic	kman line care	
	 Alteplase (Cathflo) 2 mg as needed for occluded hours, instill a 2nd dose if occluded 	catheter. For clearing central line cathe	ter- retain in catheter for 30 minutes to 2
Emergeno	ry Medications: (May give emergency medications IM if I	V route unavailable)	
	DiphenhydrAMINE (BENADRYL) 25 to 50 mg IV as nee	ded for mild to moderate drug reaction	s (flushing dizziness headaches
	oresis, fever, palpitations, chest discomfort, blood pressu once, if reaction does not resolve in 3 minutes may repe	re changes (>/= 20 points in SBP), naus	sea, urticaria, chills, pruritic). Administer 25
■ Avoid	Epinephrine 0.3 mg IM once for anaphylaxis. If reaction use of hand, foot, leg veins in elderly patient and those was a second or the contract of the contract	•	- 0
•			
Patient name:		Provider printed name:	
DOB:		Provider signature:	
Height	Weight	Date:	