



Riverbend Medical Center Regional Infusion Center (RIC)  
3377 Riverbend Drive Suite 502/510  
Springfield, Oregon 97477  
Phone 541-222-6280 Fax 541-349-8006

### Anticoagulation Bridging Order Set (v. 03/07/2024)

Diagnosis/Indication (ICD-10): \_\_\_\_\_

Follow up anti-coagulation by \_\_\_\_\_ (provider required) & \_\_\_\_\_ Anticoagulation Clinic (if applicable) after discharge from RIC

.. Bridging to warfarin (new start)

- Start warfarin plus enoxaparin bridging tomorrow (or day after hospital discharge) per pharmacy dosing

.. Bridging off/ on for procedure

Date of Procedure: \_\_\_\_\_ Type of procedure: \_\_\_\_\_

Take last dose of warfarin 5 days before procedure, or on \_\_\_\_\_

Begin enoxaparin per pharmacy dosing 4 days before procedure, or on \_\_\_\_\_

.. On day after procedure, resume warfarin plus enoxaparin bridging per pharmacy dosing (or day after hospital discharge)

.. Patient will NOT return to RIC for bridging after procedure

■ Discontinuation anticoagulation bridging order set when criteria met

- Mechanical MVR, Mechanical AVR plus a-fib: overlap for at least 4 days & INR 2.5- 3.5 for at least 1 day
- PE, DVT (new or history): overlap for at least 5 days & INR 2-3 for at least 2 days
- All other indications: overlap for at least 4 days & INR 2-3 for at least 1 day

#### Labs:

- INR, CBC no differential, BMP on day 1 (if not done in last 7 days), then per pharmacy protocol

#### Nursing communications:

- Vital signs: as needed

#### Emergency Medications:

- Diphenhydramine (BENADRYL) 25 to 50 mg IV as needed for mild to moderate drug reactions (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes ( $\geq$  20 points in SBP), nausea, urticaria, chills, pruritic). Administer 25 mg IV once, if reaction does not resolve in 3 minutes may repeat 25 mg IV dose for a total of 50 mg and contact provider.
- MethylPREDNISolone sodium succinate (Solu-MEDROL) 125 mg IV once as needed for shortness of breath, continued symptoms of mild to moderate drug reaction (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes ( $\geq$  20 points in SBP), nausea, urticaria, chills, pruritic) that worsen or persist 5 minutes after administration of diphenhydramine (Benadryl). Contact provider if given.
- Epinephrine 0.3 mg IM once for anaphylaxis. If reaction does not resolve in 3 minutes may repeat 0.3 mg IM dose for a total of 0.6 mg. Avoid use of hand, foot, leg veins in elderly patient and those with occlusive vascular disease. Contact provider if given.
- Famotidine (PEPCID) 20 mg IV once as needed for infusion/ allergic reaction.

Patient name and DOB \_\_\_\_\_

Provider printed name: \_\_\_\_\_

Provider signature: \_\_\_\_\_

Height \_\_\_\_\_ Weight \_\_\_\_\_

Date: \_\_\_\_\_ Time: \_\_\_\_\_