



Change order details by crossing out unwanted information and writing in desired details/instructions. Place a line through the **x** to remove the pre-checked option.

BENRALIZUMAB (FASENRA) THERAPY PLAN [11500850] Columbia Network Infusion Centers

Select Location: Southwest/Vancouver St John/Longview Start Date: _____

Diagnosis/Indication: _____

ICD-10 Code(s): _____ Authorization Number: _____

Patient Vitals: Height _____ Weight _____

Provider Communication Orders

To reduce delays in treatment and phone calls to your office, you may participate in the PeaceHealth formulary process by selecting this option. A clinic pharmacist will adjust orders according to PeaceHealth site approved policies and procedures.

- I agree to utilize the PeaceHealth policies & procedures that have been reviewed by the Pharmacy & Therapeutics Committee and authorized by the Medical Executive Committee of PeaceHealth. This agreement will be issued for the duration of active orders contained within this treatment plan.**

Supportive Care

Interval

- | | | | |
|-----------------------|---|---------------|---------------------|
| <input type="radio"/> | benralizumab (FASENRA) 30 mg/mL inj 30 mg x 3 doses | Every 28 days | Route: Subcutaneous |
| <input type="radio"/> | benralizumab (FASENRA) 30 mg/mL injection 30 mg | Every 56 days | Route: Subcutaneous |

Emergency Medications

Interval

- | | | | |
|-------------------------------------|--|-----|----------------------|
| <input checked="" type="checkbox"/> | diphenhydrAMINE (BENADRYL) injection 25-50 mg
<i>25-50 mg Once As Needed Intramuscular, For mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>= 20 points in SBP), nausea, urticaria, chills, pruritis). Administer 50 mg IM if patient has NOT had diphenhydramine within 2 hours of reaction. Administer 25 mg IM if patient has had diphenhydramine within 2 hours of reaction, if reaction doesn't resolve in 3 minutes may repeat 25 mg IM dose for a total of 50 mg, and notify provider.</i> | PRN | Route: Intramuscular |
| <input checked="" type="checkbox"/> | albuterol 90 mcg/actuation inhaler 2 puff
<i>2 puff Once As Needed Inhalation, Wheezing, Shortness of Breath, wheezing, shortness of breath associated with infusion reaction and notify provider. Administer with a spacer if available.</i> | PRN | Route: Inhalation |
| <input checked="" type="checkbox"/> | methylPREDNISolone sod suc(PF) (Solu-MEDROL) Injection 125 mg
<i>125 mg Once As Needed Intramuscular, For shortness of breath for continued symptoms of mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>= 20 points in SBP), nausea, urticaria, chills, pruritis) that worsen or persist 5 minutes after administration of diphenhydramine (Benadryl), and notify provider, Do not inject into deltoid.</i> | PRN | Route: Intramuscular |
| <input checked="" type="checkbox"/> | EPINEPHrine (ADRENALIN) injection 0.5 mg
<i>0.5 mg Once As Needed Intramuscular, For severe drug reaction (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort plus blood pressure changes (>= 40 points in SBP), shortness of breath with wheezing and O2Sat <90%), and notify provider, For 1 dose.</i> | PRN | Route: Intramuscular |

Provider Signature

Date

Time

Provider's Printed Name:

Place Patient Label Here

Page 1 of 1

EHR5003-DT (12/08/2024)

BENRALIZUMAB (FASENRA) THERAPY PLAN