



Change order details by crossing out unwanted information and writing in desired details/instructions. Place a line through the x to remove the pre-checked option.

IRON - FERRIC CARBOXYMALTOSE (INJECTAFER) IVPB [11500556] Columbia Network Infusion Centers

Select Location: Southwest/Vancouver St John/Longview Start Date: _____

Diagnosis/Indication: _____

ICD-10 Code(s): _____ Authorization Number: _____

Patient Vitals: Height _____ Weight _____

Provider Communication Orders

To reduce delays in treatment and phone calls to your office, you may participate in the PeaceHealth formulary process by selecting this option. A clinic pharmacist will adjust orders according to PeaceHealth site approved policies and procedures.

- I agree to utilize the PeaceHealth policies & procedures that have been reviewed by the Pharmacy & Therapeutics Committee and authorized by the Medical Executive Committee of PeaceHealth. This agreement will be issued for the duration of active orders contained within this treatment plan.**

Labs	Interval
<input checked="" type="checkbox"/> Physician Communication <i>Provider to send most recent Ferritin and/or Iron Deficiency Panel Lab results with completed order.</i>	Once

Supportive Care	Interval
<input checked="" type="checkbox"/> FERRIC CARBOXYMALTOSE IVPB 750 mg IV every 7 days x 2 doses Route: IV <i>Monitor patient for signs and symptoms of hypersensitivity DURING INFUSION AND FOR AT LEAST 30 MINUTES AFTER INFUSION. Hypersensitivity symptoms may include: anaphylaxis, flushing, dyspnea, tachycardia, and increased blood pressure. For patients weighing less than 50 kg, the dose will be updated to 15 mg/kg.</i>	
<input checked="" type="checkbox"/> SODIUM CHLORIDE 0.9 % IV BOLUS PRN Route: IV <i>IV site discomfort May use 250 ml NS at 100ml/hr. Run concurrent with iron infusions as needed.</i>	

Nursing Orders	Interval
<input checked="" type="checkbox"/> Nursing Communication <i>Monitor patient for signs and symptoms of hypersensitivity DURING INFUSION AND FOR AT LEAST 30 MINUTES AFTER INFUSION. Hypersensitivity symptoms may include: anaphylaxis, flushing, dyspnea, tachycardia, and increased blood pressure.</i>	Every 7 days
<input checked="" type="checkbox"/> Nursing Communication <i>Discontinue therapy plan when treatment complete.</i>	Every visit

Vascular Access (single select)	Interval
<input type="radio"/> IV - Insert peripheral IV - sodium chloride (NS) flush 10 mL <i>10 mL As Needed for Line Care</i>	Every Visit
<input type="radio"/> Central line (non-PICC) - Access vascular device and confirm patency	Every Visit

Provider Signature Date Time

Provider's Printed Name: _____

Place Patient Label Here

- Initiate Central line (non-PICC) maintenance protocol
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care prior to medication administration PRN
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care post medication administration PRN
 - sodium chloride (NS) flush 20 mL
20 mL As Needed for Line Care post lab draw PRN
 - sodium chloride (NS) flush 10 mL
10 mL for Line Care at discharge and de-access every visit
 - heparin 100 units/mL 5 mL
5 mL for de-access PRN
 - alteplase (CATHFLO) injection 2 mg
Other, Starting when released, For 2 dose, PRN, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded
- O** PICC line Every Visit
- Access vascular device and confirm patency
 - Initiate PICC maintenance protocol
 - Nursing Communication
Change PICC line dressing weekly and PRN
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care prior to medication administration PRN
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care post medication administration PRN
 - sodium chloride (NS) flush 20 mL
20 mL As Needed for Line Care post lab draw PRN
 - alteplase (CATHFLO) injection 2 mg
Other, Starting when released, For 2 dose, PRN, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded

PRN Medications	Interval	
<input type="radio"/> sodium chloride (NS) flush 10 mL 10 mL As Needed for Line Care	PRN	Route: IV
<input type="radio"/> sodium chloride 0.9 % bolus 500 mL 500 mL Once for Line Care, Starting when released, For 1 dose, Run at TKO (25ml/hr)	PRN	Route: IV

Emergency Medications	Interval	
<input checked="" type="checkbox"/> albuterol 90 mcg/actuation inhaler 2 puff 2 puff Once As Needed Inhalation, Wheezing, Shortness of Breath, wheezing, shortness of breath associated with infusion reaction and notify provider. Administer with a spacer if available.	PRN	Route: Inhalation

Provider Signature Date Time

Provider's Printed Name:

- X methylPREDNISolone sodium succinate (Solu-MEDROL) Injection 125 mg PRN Route: IV
125 mg Once As Needed IV, For shortness of breath for continued symptoms of mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>= 20 points in SBP), nausea, urticaria, chills, pruritis) that worsen or persist, and notify provider.

- X EPINEPHrine (ADRENALIN) injection for Allergic Reaction 0.5 mg PRN Route: Intramuscular
0.5 mg Once As Needed Intramuscular, For severe drug reaction (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort plus blood pressure changes (>= 40 points in SBP), shortness of breath with wheezing and O2Sat <90%), and notify provider.

Provider Signature

Date

Time

Provider's Printed Name: DD

Place Patient Label Here

Page 3 of 3

EHR5002-DT (12/19/2024)

IRON - FERRIC CARBOXYMALTOSE (INJECTAFER) IVPB