

## ZOLEDRONIC ACID (ZOMETA) (NOT FOR RECLAST) [11500326] Columbia Network Infusion Centers

			. 🗕		
Select Lo	ocation: O Southwest/Vancouver C	St John/Longview	Start Date:		
Diagnos	is/Indication:				
ICD-10 (	Code(s): Author	orization Number:			
Patient \	/itals: Height Weight				
Provi	der Communication Orders				
proces	luce delays in treatment and phone calls to yess by selecting this option. A clinic pharmaces and procedures.		·		
х	I agree to utilize the PeaceHealth policies of Therapeutics Committee and authorized by agreement will be issued for the duration of	the Medical Executive Co	mmittee of PeaceHealth. This		
Labs		Interval			
X	Comprehensive Metabolic Panel		ults not provided by MD office; must ithin 7 DAYS		
Supp	ortive Care	Interval			
X	ZOLEDRONIC ACID (ZOMETA) IV		Route: IV		
If creatinine clearance is greater than 60ml/min administer 4mg dose. If creatinine clearance is between 50-60ml/min administer 3.5mg dose. If creatinine clearance is between 40-49ml/min administer 3.3mg dos If creatinine clearance is between 30-39ml/min administer 3mg dose. DO NOT ADMINISTER IF CREATING CLEARANCE IS LESS THAN 30 ML/MIN.  PHARMACY TO UPDATE DOSE BASED ON CREATININE CLEARANCE.					
	Dose: _4mg_ Frequency:				
Nursi	ng Orders	Interval	Interval		
X	Nursing Communication  Treatment will be withheld for patients with	Every Visit creatinine clearance less that	an 30 mL/min.		
Vascu	ular Access (single select)	Interval			
Ο	IV - Insert peripheral IV - sodium chloride (NS) flush 10 mL 10 mL As Needed for Line Care	Every Visit			
Ο	Central line (non-PICC)  - Access vascular device and confirm par  - Initiate Central line (non-PICC) mainten  - sodium chloride (NS) flush 10 mL  10 mL As Needed for Line Care price	ance protocol	on PRN		
7	Provider Signature	Date	Time		
Provider	's Printed Name:				
	Place Patient Label Here	Page 1 of 2	EHR5064-DT (06/23/2025)		

**ZOLEDRONIC ACID (ZOMETA) (NOT FOR RECLAST)** 

- sodium chloride (NS) flush 10 mL

10 mL As Needed for Line Care post medication administration PRN

- sodium chloride (NS) flush 20 mL

20 mL As Needed for Line Care post lab draw PRN

- sodium chloride (NS) flush 10 mL

10 mL for Line Care at discharge and de-access every visit

heparin 100 units/mL 5 mL

5 mL for de-access PRN

- O PICC line Every Visit
  - Access vascular device and confirm patency
  - Initiate PICC maintenance protocol

- Nursing Communication

Change PICC line dressing weekly and PRN

- sodium chloride (NS) flush 10 mL

10 mL As Needed for Line Care prior to medication administration PRN

sodium chloride (NS) flush 10 mL

10 mL As Needed for Line Care post medication administration PRN

- sodium chloride (NS) flush 20 mL

20 mL As Needed for Line Care post lab draw PRN

- alteplase (CATHFLO) injection 2 mg

Other, Starting when released, For 2 dose, PRN, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded

PRN Medications		Interval		
О	sodium chloride (NS) flush 10 mL	PRN	Route: IV	
	10 mL As Needed for Line Care			
O	sodium chloride 0.9 % bolus 500 mL	PRN	Route: IV	
500 mL Once for Line Care, Starting when released, For 1 dose, Run at TKO (25ml/hr)				

Provider Signature	Date	Time	

Provider's Printed Name: