

## Peace Harbor Medical Center Outpatient Infusion Service Request

Phone: (541) 902-6019 Fax: (541) 902-1649

Thank you for selecting our infusion services team to care for your patient. Please provide ALL information listed below to ensure that we can process orders and schedule your patient for treatment without delay.

## Part A- Patient scheduling and contact information: Patient Name (Last, First): \_\_\_\_\_\_\_Date of Birth: \_\_\_\_\_\_ Patient Contact Information and Phone Number (s): Ordering Provider Name (Print): Provider Clinic or Service Address: Clinic or Service Phone Number: \_\_\_\_\_\_ Clinic or Service Fax Number: \_\_\_\_\_\_ Diagnosis (include ICD 10 codes): Medication and Service Requested- list J-Code/ CPT code if known: \_\_\_\_\_\_ Date Service is Requested to Begin: \_\_\_\_\_\_ Date Service is Expected to End: \_\_\_\_ Order will expire 1 year from date of provider signature unless "date service is expected to end" is earlier. Part B- Insurance and Prior Authorization. Any non-PeaceHealth provider must obtain prior authorization prior to service. Attach a copy of authorization documentation received from insurance payer when submitting orders. Insurance (Payer) Company: Prior Authorization Number and Conditions: Prior Authorization Expiration Date: Insurance (Payer) Contact Phone Number: Part C- Elements needed to guide medication therapy are included with request for service: Orders and instruction (use the PeaceHealth approved ordering form if you are not a PeaceHealth provider) are complete and include provider signature at the bottom of each page. Check the boxes of ALL orders you would like to activate. For blood products, PeaceHealth Blood and Transfusion Consent form is signed and dated by the provider and the patient. If information is located outside of PeaceHealth's electronic medical record system attach the following: A list of current medications reconciled by patient provider is available and includes a list of known allergies. Recent progress notes from ordering provider. A copy of relevant laboratory results and other appropriate supporting documentation. IMPORTANT MESSAGE TO PROVIDERS: To reduce delays in treatment and added phone calls, you may participate to utilize PeaceHealth preferred medication formulary options by signing this document. A clinical pharmacist will adjust orders to align with PeaceHealth medical staff approved medication formulary options, policies, and procedures. I agree to utilize PeaceHealth preferred medication formulary options, policies and procedures that have been authorized by PeaceHealth Medical Staff. This agreement will be issued for the duration of active orders contained within this treatment plan. PROVIDER SIGNATURE: \_\_\_\_\_ DATE: \_\_\_\_\_ TIME:\_\_\_\_

FAX completed service request and orders to: PHMC OP Infusion and Nursing Services 541-902-1649



Progress & Orders



## MethylPREDNISolone (SoluMedrol) **Outpatient Infusion Therapy Plan**

	ders Are Initiated by Default Unless Crossed Out by Practitioner. All <u>Boxed Orders</u> Require Practitioner Check to be Initiated.		
Heading Supportive Care	Content  MathylppppNICalana and ium aussinata (Calu Madral) IV infraian.		
Supportive care			
	Select Dose:		
	☐ 500 mg		
	☐ 1000 mg		
	mg		
	Select Frequency:		
	☐ Once (single dose)		
	☐ Once daily for 2 days		
	☐ Once daily for 3 days		
	☐ Once daily for 5 days		
	Select Interval:		
	Repeat every 14 days		
	Repeat every 28 days		
Nursing IV Access	Select the most appropriate option below:		
and Maintenance			
	Sodium chloride 0.9% (NS) flush 10 mL IV once as needed for line care.		
	Za a de la manta e di son (1.10) masin 10 mili me da le me de la me da le m		
	☐ Access and use NON-PICC Central Line/CVAD		
	☐ Initiate Central Line (non-PICC) maintenance protocol.		
	☑ Sodium chloride 0.9% (NS) flush 10 mL IV as needed for line care, before and after medication		
	administration, at discharge, and at de-access (sterile NS for Port-a-Cath access)		
	☑ Sodium chloride 0.9% (NS) flush 20 mL IV as needed for line care post lab draw.		
	☐ Heparin, porcine (PF) 100 unit/mL flush 5 mL IV as needed for line care, for de-access.		
	☐ Alteplase (Cathflo) injection 2 mg intra-catheter as needed for occluded central line catheters-		
	For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand		
	undisturbed to allow large bubbles to dissipate. Mix by gently swirling until completely		
	dissolved (complete dissolution should occur within 3 minutes); do not shake. Final		
	concentration: 1mg/mL. Instill medication in non-functional lumen. Do not use lumen while		
	dwelling. Allow to dwell 30 minutes and check for patency by drawing back on lumen for blood		
	return. If line is still not patent, allow medication to dwell an additional 90 minutes. Dwell time		
	not to exceed 120 min. Use second dose of Alteplase (Cathflo) if catheter is not patent after 120		
	min. If the catheter is functional, aspirate and waste the medication and residual clot prior to		
	flushing the line.		
	☐ Access and use PICC Central Line/CVAD		
	☐ Change PICC line dressing weekly and as needed.		
	☑ Sodium chloride 0.9% (NS) flush 10 mL IV as needed for line care, and before and after		
	medication administration.		
	☑ Sodium chloride 0.9% (NS) flush 20 mL IV as needed for line care post lab draw		
	☐ Alteplase (Cathflo) injection 2 mg intra-catheter as needed for occluded central line catheters-		
	For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand		

Practitioner Signature: \_\_

Final page of orders must include signature of the ordering practitioner, date, and time.



Progress & Orders



## MethylPREDNISolone (SoluMedrol) Outpatient Infusion Therapy Plan

All Pre-Selected Boxed Orders Are Initiated by Default Unless Crossed Out by Practitioner. All Boxed Orders Require Practitioner Check to be Initiated.

Heading	Content		
	undisturbed to allow large bubbles to dissipate. Mix by gently swirling until completely		
	dissolved (complete dissolution should occur within 3 minutes); do not shake. Final		
	concentration: 1mg/mL. Instill medication in non-functional lumen. Do not use lumen while		
	dwelling. Allow to dwell 30 minutes and check for patency by drawing back on lumen for blood		
	return. If line is still not patent, allow medication to dwell an additional 90 minutes. Dwell time		
	not to exceed 120 min. Use second dose of Alteplase (Cathflo) if catheter is not patent after 120		
	min. If the catheter is functional, aspirate and waste the medication and residual clot prior to		
	flushing the line.		
As Needed	Standard As Needed Medications:		
Medications	Sodium chloride 0.9% (NS) flush 10 mL IV as needed for line care.		
	☑ Sodium chloride 0.9% 500 mL continuous infusion at 25 mL/hour IV as needed for therapy		
	administration (i.e., blood products, chemotherapy, potassium administration).		
Referral			
PHMC Outpatient	PROVIDER – PLEASE SIGN, DATE AND TIME ORDERS AND RETURN TO:		
Infusion Contact	PeaceHealth Peace Harbor Medical Center Outpatient Infusion Services Department		
Information	400 Ninth Street, Florence, OR 97439		
	Contact Phone: 541-902-6019 and FAX <b>541-902-1649</b>		
Authorization by	Person giving verbal or telephone order:		
Verbal or	Person receiving verbal or telephone order:		
Telephone Order	☐ Check to indicate verbal or telephone orders have been read back to confirm accuracy		

Practitioner Signature:	Date of Or	der:Time:

Final page of orders must include signature of the ordering practitioner, date, and time.