



Change order details by crossing out unwanted information and writing in desired details/instructions.
Place a line through the x to remove the pre-checked option.

INFliximab OR BIOSIMILAR MAINTENANCE INFUSION [11500956] Columbia Network Infusion Centers

Select Location: Southwest/Vancouver St John/Longview Start Date: _____

Diagnosis/Indication: _____

ICD-10 Code(s): _____ Authorization Number: _____

Patient Vitals: Height _____ Weight _____

Provider Communication Orders

To reduce delays in treatment and phone calls to your office, you may participate in the PeaceHealth formulary process by selecting this option. A clinic pharmacist will adjust orders according to PeaceHealth site approved policies and procedures.

I agree to utilize the PeaceHealth policies & procedures that have been reviewed by the Pharmacy & Therapeutics Committee and authorized by the Medical Executive Committee of PeaceHealth. This agreement will be issued for the duration of active orders contained within this treatment plan.

Pre-Screening Labs

Provider Communication
Provider to send PPD/Tuberculosis screening and Hep B screening lab results with completed orders.

Labs	Interval
<input type="radio"/> Complete Blood Count W/ Automated Differential	Frequency _____
<input type="radio"/> Comprehensive Metabolic Panel	Frequency _____

Pre-Medications	Interval
<input type="radio"/> acetaminophen (TYLENOL) tablet 650 mg	Every Visit Route: Oral
<input type="radio"/> loratadine (CLARITIN) tablet 10 mg	Every Visit Route: Oral
<input type="radio"/> diphenhydramine (BENADRYL) capsule 25 mg	Every Visit Route: Oral
<input type="radio"/> methylPREDNISolone sodium succinate (Solu-MEDROL) Injection 40 mg	Every Visit Route: IV

Supportive Care	Interval
PLEASE SELECT (AND OBTAIN INSURANCE AUTHORIZATION IF REQUIRED) FOR ONE OF THE PEACEHEALTH PREFERRED AGENTS BELOW. THEY ARE LISTED IN ORDER OF PREFERENCE.	
<input type="radio"/> INFliximab IVPB (REMICADE) <i>Infuse 10ml/hr x15min; then 20ml/hr x15min; then 40ml/hr x15min; then 80ml/hr x15min; then 150ml/hr x15min; then max 250ml/hr until done.</i>	Route: IV Dose _____ Frequency _____
<input type="radio"/> INFliximab-ABDA IVPB (RENFLEXIS) <i>Infuse 10ml/hr x15min; then 20ml/hr x15min; then 40ml/hr x15min; then 80ml/hr x15min; then 150ml/hr x15min; then max 250ml/hr until done.</i>	Route: IV Dose _____ Frequency _____
<input type="radio"/> INFliximab-DYYB IVPB (INFLECTRA) <i>Infuse 10ml/hr x15min; then 20ml/hr x15min; then 40ml/hr x15min; then 80ml/hr x15min; then 150ml/hr x15min; then max 250ml/hr until done.</i>	Route: IV Dose _____ Frequency _____

Provider Signature _____ Date _____ Time _____

Provider's Printed Name: _____

Nursing Orders		Interval
<input checked="" type="checkbox"/> Nursing Communication		Every Visit <i>Assess patients' vital signs prior to the infliximab infusion, before each rate increase, and 30 minutes following the infusion. Observe patient 30 minutes after completion of infusion.</i>
Vascular Access (single select)		
<input type="radio"/> IV		Every Visit
<ul style="list-style-type: none"> - Insert peripheral IV - sodium chloride (NS) flush 10 mL <i>10 mL As Needed for Line Care</i> 		
<input type="radio"/> Central line (non-PICC)		Every Visit
<ul style="list-style-type: none"> - Access vascular device and confirm patency - Initiate Central line (non-PICC) maintenance protocol - sodium chloride (NS) flush 10 mL <i>10 mL As Needed for Line Care prior to medication administration PRN</i> - sodium chloride (NS) flush 10 mL <i>10 mL As Needed for Line Care post medication administration PRN</i> - sodium chloride (NS) flush 20 mL <i>20 mL As Needed for Line Care post lab draw PRN</i> - sodium chloride (NS) flush 10 mL <i>10 mL for Line Care at discharge and de-access every visit</i> - heparin 100 units/mL 5 mL <i>5 mL for de-access PRN</i> - alteplase (CATHFLO) injection 2 mg <i>Other, Starting when released, For 2 dose, PRN, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded</i> 		
<input type="radio"/> PICC line		Every Visit
<ul style="list-style-type: none"> - Access vascular device and confirm patency - Initiate PICC maintenance protocol - Nursing Communication <i>Change PICC line dressing weekly and PRN</i> - sodium chloride (NS) flush 10 mL <i>10 mL As Needed for Line Care prior to medication administration PRN</i> - sodium chloride (NS) flush 10 mL <i>10 mL As Needed for Line Care post medication administration PRN</i> - sodium chloride (NS) flush 20 mL <i>20 mL As Needed for Line Care post lab draw PRN</i> - alteplase (CATHFLO) injection 2 mg <i>Other, Starting when released, For 2 dose, PRN, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded</i> 		

Provider Signature

Date

Time

Provider's Printed Name:

Place Patient Label Here

Page 2 of 3

EHR5029-DT (02/03/2026)

INFILIXIMAB OR BIOSIMILAR MAINTENANCE INFUSION

PRN Medications		Interval
<input type="radio"/> sodium chloride (NS) flush 10 mL <i>10 mL As Needed for Line Care</i>	PRN	Route: IV
<input type="radio"/> sodium chloride 0.9 % bolus 500 mL <i>500 mL Once for Line Care, Starting when released, For 1 dose, Run at TKO (25ml/hr)</i>	PRN	Route: IV
Emergency Medications		Interval
<input checked="" type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25-50 mg <i>25-50 mg Once As Needed IV, For mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>= 20 points in SBP), nausea, urticaria, chills, pruritis). Administer 50 mg IV if patient has NOT had diphenhydramine within 2 hours of reaction. Administer 25 mg IV if patient has had diphenhydramine within 2 hours of reaction, if reaction doesn't resolve in 3 minutes may repeat 25 mg IV dose for a total of 50 mg, and contact provider.</i>	PRN	Route: IV
<input checked="" type="checkbox"/> albuterol 90 mcg/actuation inhaler 2 puff <i>2 puff Once As Needed Inhalation, Wheezing, Shortness of Breath, wheezing, shortness of breath associated with infusion reaction and notify provider. Administer with a spacer if available.</i>	PRN	Route: Inhalation
<input checked="" type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) Injection 125 mg <i>125 mg Once As Needed IV, For shortness of breath for continued symptoms of mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>= 20 points in SBP), nausea, urticaria, chills, pruritis) that worsen or persist 5 minutes after administration of diphenhydramine (Benadryl), and contact provider.</i>	PRN	Route: IV
<input checked="" type="checkbox"/> EPINEPHRine (ADRENALIN) injection for Allergic Reaction 0.5 mg <i>0.5 mg Once As Needed Intramuscular, For severe drug reaction (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort plus blood pressure changes (>= 40 points in SBP), shortness of breath with wheezing and O2Sat <90%), and contact provider.</i>	PRN	Route: Intramuscular

Provider Signature

Date

Time

Provider's Printed Name:

Place Patient Label Here

Page 3 of 3

EHR5029-DT (02/03/2026)

INFLIXIMAB OR BIOSIMILAR MAINTENANCE INFUSION