



Thank you for selecting our infusion services team to care for your patient. Please provide ALL information listed below to ensure that we can process orders and schedule your patient for treatment without delay.

Part A- Patient scheduling and contact information:

Patient Name (Last, First): _____ Date of Birth: _____

Patient Contact Information and Phone Number (s): _____

Ordering Provider Name (Print): _____

Provider Clinic or Service Address: _____

Clinic or Service Phone Number: _____ Clinic or Service Fax Number: _____

Diagnosis (include ICD 10 codes): _____

Medication and Service Requested- list J-Code/ CPT code if known: _____

Date Service is Requested to Begin: _____ **Date Service is Expected to End:** _____

Order will expire 1 year from date of provider signature unless "date service is expected to end" is earlier.

Part B- Insurance and Prior Authorization. Any non-PeaceHealth provider must obtain prior authorization prior to service. Attach a copy of authorization documentation received from insurance payer when submitting orders.

Insurance (Payer) Company: _____

Prior Authorization Number and Conditions: _____

Prior Authorization Expiration Date: _____

Insurance (Payer) Contact Phone Number: _____

Part C- Elements needed to guide medication therapy are included with request for service:

- Orders and instruction (use the PeaceHealth approved ordering form if you are not a PeaceHealth provider) are complete and include provider signature at the bottom of each page. Check the boxes of ALL orders you would like to activate.
- For blood products, PeaceHealth Blood and Transfusion Consent form is signed and dated by the provider and the patient.

If information is located outside of PeaceHealth's electronic medical record system attach the following:

- A list of current medications reconciled by patient provider is available and includes a list of known allergies.
- Recent progress notes from ordering provider.
- A copy of relevant laboratory results and other appropriate supporting documentation.

IMPORTANT MESSAGE TO PROVIDERS: To reduce delays in treatment and added phone calls, you may participate to utilize PeaceHealth preferred medication formulary options by signing this document. A clinical pharmacist will adjust orders to align with PeaceHealth medical staff approved medication formulary options, policies, and procedures.

I agree to utilize PeaceHealth preferred medication formulary options, policies and procedures that have been authorized by PeaceHealth Medical Staff. This agreement will be issued for the duration of active orders contained within this treatment plan.

PROVIDER SIGNATURE: _____ **DATE:** _____ **TIME:** _____

FAX completed service request and orders to: PHMC OP Infusion and Nursing Services 541-902-1649



Ferric Carboxymaltose (Injectafer) Outpatient Infusion Therapy Plan

All Pre-Selected Boxed Orders Are Initiated by Default Unless Crossed Out by Practitioner. All Boxed Orders Require Practitioner Check to be Initiated.

| Heading | Content |
|---------------------------------|---|
| For Admission to Service | <p>Provider Instruction – Please review information below and address requirements for admission to service:</p> <ol style="list-style-type: none"> 1. Provider to order CBC with automated differential and iron deficiency panel prior to therapy 2. Not FDA-approved for iron deficiency anemia in dialysis patients |
| Supportive Care | <p>Choose one of the following (A-B):</p> <p>A. Less than 50 kg: <input type="checkbox"/> Ferric carboxymaltose (Injectafer) 15 mg/kg IV infused over 20 minutes every 7 days x 2 doses</p> <p>B. Greater than or equal to 50 kg: <input type="checkbox"/> Ferric carboxymaltose (Injectafer) 750 mg IV infused over 20 minutes every 7 days x 2 doses</p> <p>C. <input checked="" type="checkbox"/> Sodium chloride 0.9% (NS) continuous infusion at 100 mL/hour IV as needed for IV site discomfort. Run concurrent with iron infusions as needed.</p> |
| Nursing Orders | <p><input checked="" type="checkbox"/> Monitor patient for signs and symptoms of hypersensitivity during infusion and for at least 30 minutes after infusion. Hypersensitivity symptoms may include anaphylaxis, flushing, dyspnea, tachycardia, and increased blood pressure.</p> <p><input checked="" type="checkbox"/> Monitor patient for MILD hypersensitivity reactions and initiate interventions as indicated: Mild Hypersensitivity Symptoms: Metallic taste, chest tightening, back tightening, increased anxiety, flushing (normothermic), dizziness, nausea, headache, diaphoresis (normothermic), palpitations, fish bone reaction (transient flushing, chest and back tightening, joint pain).</p> <p>Step 1: STOP infusion, monitor vital signs and O2 every 5 minutes (until symptoms resolve or for a minimum of 15 minutes).</p> <p>Step 2: Administer Methylprednisolone as ordered.</p> <p>Step 3: Resume infusion at reduced rate of 50% original rate.</p> <p>Step 4: Observe for 1 hour following infusion completion.</p> <p>Step 5: Notify provider of adverse reaction.</p> <p><input checked="" type="checkbox"/> Monitor patient for MODERATE hypersensitivity reactions and initiate intervention as indicated: Moderate Hypersensitivity Symptoms: Mild symptoms PLUS chest discomfort (pressure), shortness of breath, hypo/hypertension (greater than or equal to a 20-point change in SBP), increased temperature (greater than or equal to 38 C or 100.4 F) with rigors, urticaria, new onset or progressive edema</p> <p><input checked="" type="checkbox"/> Monitor for ANAPHYLAXIS: Mild and moderate symptoms PLUS hypo/hypertension (greater than or equal to a 40-point change in SBP), shortness of breath (RR greater than 20, O2 sat less than 90%) with wheezing or stridor, swelling of face/tongue, corneal edema, or periorbital edema.</p> <p>Step 1: STOP infusion and activate rapid response.</p> <p>Step 2: Monitor vital signs and O2 sat every 5 minutes, place patient in left lateral tilt (if pregnant).</p> <p>Step 3: Administer Methylprednisolone as ordered.</p> <p>Step 4: Administer Famotidine as ordered.</p> |

Practitioner Signature: _____ **Date of Order:** _____ **Time:** _____

Final page of orders must include signature of the ordering practitioner, date, and time.



Ferric Carboxymaltose (Injectafer) Outpatient Infusion Therapy Plan

All Pre-Selected Boxed Orders Are Initiated by Default Unless Crossed Out by Practitioner. All Boxed Orders Require Practitioner Check to be Initiated.

| Heading | Content |
|--|--|
| | Step 5: Administer Epinephrine as ordered (ONLY FOR ANAPHYLAXIS). Step 6: Transfer patient to main Emergency Department as soon as possible. |
| Nursing IV Access and Maintenance | <p>Select the most appropriate option below:</p> <ul style="list-style-type: none"> <input checked="" type="checkbox"/> Insert PERIPHERAL IV once as needed <ul style="list-style-type: none"> <input checked="" type="checkbox"/> Sodium chloride 0.9% (NS) flush 10 mL IV once as needed for line care <input type="checkbox"/> Access and use NON-PICC Central Line/CVAD as needed and confirm patency <ul style="list-style-type: none"> <input checked="" type="checkbox"/> Initiate Central Line (Non-PICC) maintenance protocol <input checked="" type="checkbox"/> Sodium chloride 0.9% (NS) injection 10 mL IV as needed for line care before and after medication administration, at discharge, and at de-access <input checked="" type="checkbox"/> Sodium chloride 0.9% (NS) injection 20 mL IV once as needed for line care post lab draw <input checked="" type="checkbox"/> Heparin, porcine (PF) 100 unit/mL flush 5 mL IV as needed for line care, for de-access <input checked="" type="checkbox"/> Alteplase (Cathflo) injection 2 mg intra-catheter once as needed x 2 doses. For clearing central line catheter. Reconstitute with 2.2 mL Sterile Water to the vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix by gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Instill medication in non-functional lumen. Do not use lumen while dwelling. Allow to dwell 30 minutes and check for patency by drawing back on lumen for blood return. If line is still not patent, allow medication to dwell an additional 90 minutes. Dwell time not to exceed 120 min. Use second dose of Alteplase (Cathflo) if catheter not patent after 120 min. If the catheter is functional, aspirate and waste the medication and residual clot prior to flushing the line. <input type="checkbox"/> Access and use PICC Central Line/CVAD as needed and confirm patency <ul style="list-style-type: none"> <input checked="" type="checkbox"/> Initiate PICC maintenance protocol <input checked="" type="checkbox"/> Change PICC line dressing weekly and as needed <input checked="" type="checkbox"/> Sodium chloride 0.9% (NS) injection 10 mL IV as needed for line care before and after medication administration <input checked="" type="checkbox"/> Sodium chloride 0.9% (NS) injection 20 mL IV once as needed for line care post lab draw <input checked="" type="checkbox"/> Alteplase (Cathflo) injection 2 mg intra-catheter once as needed x 2 doses. For clearing central line catheter. Reconstitute with 2.2 mL Sterile Water to the vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix by gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Instill medication in non-functional lumen. Do not use lumen while dwelling. Allow to dwell 30 minutes and check for patency by drawing back on lumen for blood return. If line is still not patent, allow medication to dwell an additional 90 minutes. Dwell time not to exceed 120 min. Use second dose of Alteplase (Cathflo) if catheter not patent after 120 min. If the catheter is functional, aspirate and waste the medication and residual clot prior to flushing the line. |
| As Needed Medications | <p>Standard As Needed Medications:</p> <ul style="list-style-type: none"> <input checked="" type="checkbox"/> Sodium chloride 0.9% (NS) flush 10 mL IV as needed for line care <input checked="" type="checkbox"/> Sodium chloride 0.9% 500 mL continuous infusion at 25 mL/hour IV as needed for line care <input checked="" type="checkbox"/> Ondansetron 8 mg IV once as needed for nausea or vomiting. |

Practitioner Signature: _____ Date of Order: _____ Time: _____

Final page of orders must include signature of the ordering practitioner, date, and time.

Patient Identification Label



**Ferric Carboxymaltose (Injectafer)
Outpatient Infusion Therapy Plan**

All Pre-Selected Boxed Orders Are Initiated by Default Unless Crossed Out by Practitioner. All Boxed Orders Require Practitioner Check to be Initiated.

| Heading | Content |
|---|---|
| Emergency Medications | <p>If patient has symptoms of anaphylaxis (wheezing, dyspnea, hypotension, angioedema, chest pain, or tongue swelling), discontinue infusion and initiate standard emergency response procedures.</p> <p>Standard Emergency Medications:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Albuterol 90 mcg/actuation inhaler 2 puffs once as needed for wheezing, shortness of breath associated with infusion reaction and notify provider. Administer with a spacer if available. <input checked="" type="checkbox"/> MethylPREDNISolone (Solu-Medrol) injection 125 mg IV once as needed for shortness of breath for continued symptoms of mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (greater than or equal to 20 points in SBP), nausea, urticaria, chills, pruritis) that worsen or persist and notify provider. <input checked="" type="checkbox"/> Famotidine (Pepcid) injection 20 mg IV once as needed for continued symptoms of moderate hypersensitivity reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (greater than or equal to 20 points in SBP), nausea, urticaria, chills, pruritis) that worsen or persist and notify provider. <input checked="" type="checkbox"/> EPINEPHrine (Adrenalin) injection 0.5 mg IM once as needed for severe drug reaction (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort plus blood pressure changes (greater than or equal to 40 points in SBP), shortness of breath with wheezing and O2 Sat less than 90% and notify provider. |
| Referral | <input checked="" type="checkbox"/> Ambulatory referral to OP Infusion Services |
| PHMC Outpatient Infusion Contact Information | <p>PROVIDER – PLEASE SIGN, DATE AND TIME ORDERS AND RETURN TO:</p> <p>PeaceHealth Peace Harbor Medical Center Outpatient Infusion Services Department 400 Ninth Street, Florence, OR 97439 Contact Phone: 541-902-6019 and FAX 541-902-1649</p> |
| Authorization by Verbal or Telephone Order | <p>Person giving verbal or telephone order: _____</p> <p>Person receiving verbal or telephone order: _____</p> <p><input type="checkbox"/> Check to indicate verbal or telephone orders have been read back to confirm accuracy</p> |

Practitioner Signature: _____ **Date of Order:** _____ **Time:** _____

Final page of orders must include signature of the ordering practitioner, date, and time.