



Cottage Grove Infusion
1515 Village Drive
Cottage Grove, OR 97424
Phone 541-767-5447 Fax 541-767-5399 E-Fax 541-434-3164

Therapeutic Phlebotomy Order Set (v. 05/24/2024)

Diagnosis/Indication (ICD-10): _____

Medications:

- .. Sodium chloride 0.9% bolus 250 mL IV over 15 minutes after phlebotomy
- .. Sodium chloride 0.9% bolus 500 mL IV over 30 minutes after phlebotomy
- .. If patient symptomatic post phlebotomy, sodium chloride 0.9% bolus 500 mL IV over 30 minutes pm lightheaded or dizzy

Nursing communications:

- n Vital signs: before, after, and as needed during procedure
- n Therapeutic phlebotomy: Volume remove _____ Frequency: _____
- n Draw CBC if there are no results available within previous 7 days
- .. Serial patient: hold phlebotomy until next scheduled appointment for hemoglobin less than _____
- .. Serial patient: phlebotomize for hemoglobin over _____

Access:

- .. Insert peripheral IV
 - Every visit, remove after IV administration complete
- .. Access & Use Central Line/ CVAD
 - Initiate Central Line (Non-PICC) Maintenance Protocol
 - Heparin, porcine (PF) 100 unit/mL flush 5 mL as needed for Port-a-Cath line care
 - Alteplase (Cathflo) 2 mg as needed for occluded catheter. For clearing central line catheter- retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded
- .. Access & Use PICC
 - Initiate PICC Maintenance Protocol
 - Normal saline flush 3 mL as needed for PICC/ Hickman line care
 - Alteplase (Cathflo) 2 mg as needed for occluded catheter. For clearing central line catheter- retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded

Emergency Medications:

- Diphenhydramine (BENADRYL) 25 to 50 mg IV as needed for mild to moderate drug reactions (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (\geq 20 points in SBP), nausea, urticaria, chills, pruritic). Administer 25 mg IV once, if reaction does not resolve in 3 minutes may repeat 25 mg IV dose for a total of 50 mg and contact provider.
- MethylPREDNISolone sodium succinate (Solu-MEDROL) 125 mg IV once as needed for shortness of breath, continued symptoms of mild to moderate drug reaction (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (\geq 20 points in SBP), nausea, urticaria, chills, pruritic) that worsen or persist 5 minutes after administration of diphenhydramine (Benadryl). Contact provider if given.
- Epinephrine 0.3 mg IM once for anaphylaxis. If reaction does not resolve in 3 minutes may repeat 0.3 mg IM dose for a total of 0.6 mg. Avoid use of hand, foot, leg veins in elderly patient and those with occlusive vascular disease. Contact provider if given.
- Famotidine (PEPCID) 20 mg IV once as needed for infusion/ allergic reaction.
- May give emergency medications IM if IV route unavailable

Patient name: _____

Provider printed name: _____

DOB: _____

Provider signature: _____

Height _____ Weight _____

Date: _____ Time: _____