PRGORD Progress & Order

PeaceHealth

Peace Harbor Medical Center
Outpatient Infusion Service Request

Phone: (541) 902-6019 Fax: (541) 902-1649

Thank you for selecting our infusion services team to care for your patient. Please provide ALL information listed below to ensure that we can process orders and schedule your patient for treatment without delay.

Part A- Patient scheduling and contact information:

Patient Name (Last, First): _____ Date of Birth: _____

Patient Contact Information and Phone Number(s): _____

Ordering Provider Name (Print): _____

Provider Clinic or Service Address: _____

Clinic or Service Phone Number: _____ Clinic or Service Fax Number: _____

Diagnosis (include ICD 10 codes): _____

Medication and Service Requested - list J-Code/CPT code if known: _____

Date Service is Requested to Begin: _____ **Date Service is Expected to End**: _____

Order will expire 1 year from date of provider signature unless "date service is expected to end" is earlier.

Part B- Insurance and Prior Authorization. Any non-PeaceHealth provider must obtain prior authorization prior to service. *Attach a copy of authorization documentation received from insurance payer when submitting orders.*

Insurance (Payer) Company: _____

Prior Authorization Number and Conditions: _____

Prior Authorization Expiration Date: _____

Insurance (Payer) Contact Phone Number: _____

Part C- Elements needed to guide medication therapy are included with request for service:

☐ Orders and instruction (use the PeaceHealth approved ordering form if you are not a PeaceHealth provider) are complete and include provider signature at the bottom of each page. Check the boxes of ALL orders you would like to activate.

☐ For blood products, PeaceHealth Blood and Transfusion Consent form is signed and dated by the provider and the patient.

If information is located outside of PeaceHealth's electronic medical record system attach the following:

☐ A list of current medications reconciled by patient provider is available and includes a list of known allergies.

☐ Recent progress notes from ordering provider.

☐ A copy of relevant laboratory results and other appropriate supporting documentation.

IMPORTANT MESSAGE TO PROVIDERS: To reduce delays in treatment and added phone calls, you may participate to utilize PeaceHealth preferred medication formulary options by signing this document. A clinical pharmacist will adjust orders to align with PeaceHealth medical staff approved medication formulary options, policies, and procedures.

I agree to utilize PeaceHealth preferred medication formulary options, policies and procedures that have been authorized by PeaceHealth Medical Staff. This agreement will be issued for the duration of active orders contained within this treatment plan.

PROVIDER SIGNATURE: _____ DATE: _____ TIME:_____

FAX completed service request and orders to: PHMC OP Infusion and Nursing Services 541-902-1649

Sensitivity: General Business Use. This document contains proprietary information and is intended for business use only.

Patient Identification Label

PeaceHealth
Peace Harbor Medical Center

Potassium Replacement
Outpatient Infusion Therapy Plan

All Pre-Selected Boxed Orders Are Initiated by Default Unless Crossed Out by Practitioner. All Boxed Orders Require Practitioner Check to be Initiated.

Heading	Content
Supportive Care	**Choose one of the following:** ☐ Potassium chloride _____ mEq IV **once** ☐ Potassium chloride IV **weekly** based on lab value. Check lab results prior to releasing this order: • For potassium serum level 3.2-3.4 mmol/L, give 20 mEq potassium chloride IV • For potassium serum level 2.5-3.1 mmol/L, give 40 mEq potassium chloride IV • For potassium serum level less than 2.5 mmol/L, start 40 mEq potassium chloride IV and call provider for additional orders. ☐ OTHER: Potassium chloride IV **_____ (frequency)** based on lab value. Check lab results prior to releasing this order: • For potassium serum level _____, give _____ mEq potassium chloride IV
Labs	☒ Serum potassium level prior to each treatment ☒ Treatment lab instructions – Provider approves to release and draw labs 2 days pre and post this planned treatment date.
Nursing Orders	☒ Nursing communication – Check serum potassium and infuse ordered dose of medication based on lab values as indicated with provider order.
Nursing IV Access and Maintenance	**Select the most appropriate option below:** ☒ **Insert PERIPHERAL IV** as needed ☒ Sodium chloride 0.9% (NS) flush 10 mL IV once as needed for line care ☐ **Access and use NON-PICC Central Line/CVAD** as needed and confirm patency ☒ Initiate Central Line (Non-PICC) maintenance protocol ☒ Sodium chloride 0.9% (NS) injection 10 mL IV as needed for line care before and after medication administration, at discharge, and at de-access ☒ Sodium chloride 0.9% (NS) injection 20 mL IV once as needed for line care post lab draw ☒ Heparin, porcine (PF) 100 unit/mL flush 5 mL IV as needed for line care, for de-access ☒ Alteplase (Cathflo) injection 2 mg intra-catheter as needed for clearing central line catheter. Reconstitute with 2.2 mL sterile water for injection to the vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix by gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Instill medication in non-functional lumen. Do not use lumen while dwelling. Allow to dwell 30 minutes and check for patency by drawing back on lumen for blood return. If line is still not patent, allow medication to dwell an additional 90 minutes. Dwell time not to exceed 120 minutes. Use second dose of Alteplase (Cathflo) if catheter not patent after 120 minutes. If the catheter is functional, aspirate and waste the medication and residual clot prior to flushing line. ☐ **Access and use PICC Central Line/CVAD** as needed and confirm patency ☒ Initiate PICC maintenance protocol ☒ Change PICC line dressing weekly and as needed ☒ Sodium chloride 0.9% (NS) injection 10 mL IV as needed for line care before and after medication administration, at discharge, and at de-access ☒ Sodium chloride 0.9% (NS) injection 20 mL IV once as needed for line care post lab draw ☒ Alteplase (Cathflo) injection 2 mg intra-catheter as needed for clearing central line catheter

Practitioner Signature: _____Date of Order: _____Time: _____

Orders must include signature of the ordering practitioner, date, and time.

Sensitivity: General Business Use. This document contains proprietary information and is intended for business use only.

Patient Identification Label

PeaceHealth
Peace Harbor Medical Center

Potassium Replacement
Outpatient Infusion Therapy Plan

All Pre-Selected Boxed Orders Are Initiated by Default Unless Crossed Out by Practitioner. All Boxed Orders Require Practitioner Check to be Initiated.

Heading	Content
Emergency Medications	**If patient has symptoms of anaphylaxis (wheezing, dyspnea, hypotension, angioedema, chest pain, or tongue swelling), discontinue infusion and initiate standard emergency response procedures.** ☒ **DiphenhydrAMINE (Benadryl)** injection 25-50 mg IV once as needed for mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (≥ 20 points in SBP), nausea, urticaria, chills, pruritis). • Administer 50 mg IV if patient has NOT had diphenhydramine within 2 hours of reaction. • Administer 25 mg IV if patient has had diphenhydramine within 2 hours of reaction, if reaction doesn't resolve in 3 minutes may repeat 25 mg IV dose for a total of 50 mg and notify provider. ☐ **Albuterol 90 mcg/actuation** inhaler 2 puffs once as needed for wheezing, shortness of breath associated with infusion reaction and notify provider. Administer with a spacer if available. ☒ **MethylPREDNISolone (Solu-Medrol)** injection 125 mg IV once as needed for shortness of breath for continued symptoms of mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (≥ 20 points in SBP), nausea, urticaria, chills, pruritis) that worsen or persist after administration of diphenhydramine (Benadryl) and notify provider. ☒ **EPINEPHrine (Adrenalin)** injection 0.5 mg IM once as needed for severe drug reaction (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort plus blood pressure changes (≥ 40 points in SBP), shortness of breath with wheezing and 02 Sat < 90%) and notify provider.
Referral	☒ Ambulatory referral to OP Infusion Services
Authorization by Verbal or Telephone Order	Person giving verbal or telephone order: _____ Person receiving verbal or telephone order: _____ ☐ Check to indicate verbal or telephone orders have been read back to confirm accuracy

Practitioner Signature: _____Date of Order: _____Time: _____

Orders must include signature of the ordering practitioner, date, and time.

Sensitivity: General Business Use. This document contains proprietary information and is intended for business use only.